

ELITE SWINE PRODUCTS

PH 1 DEN/CTC

MEDICATED

A COMPLETE PELLETTED FEED FOR NURSERY PIGS WEIGHING
FROM LESS THAN 12 LB TO 15 LB

Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline. For the control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin.

ACTIVE DRUG INGREDIENTS

Chlortetracycline	400 g/ton
Tiamulin	35 g/ton

GURANTEED ANALYSIS

Crude Protein	Not Less Than	23.5%
Lysine	Not Less Than	1.52%
Crude Fat	Not Less Than	4.0%
Crude Fiber	Not More Than	3.5%
Calcium (Ca)	Not Less Than	0.95%
Calcium (Ca)	Not More Than	1.45%
Phosphorus (P)	Not Less Than	0.69%
Salt (NaCl)	Not Less Than	0.95%
Salt (NaCl)	Not More Than	1.45%
Selenium (Se)	Not Less Than	0.29 ppm
Zinc (Zn)	Not Less Than	160 ppm

INGREDIENTS

Corn, Soybean Meal, Oats, Whey, Enzymatically Hydrolyzed Animal Protein (No Ruminant Proteins), Egg Product, Dried *Lactobacillus acidophilus* Fermentation Product, Spray Dried Blood Meal, Monocalcium Phosphate, Calcium Carbonate, , Zinc Oxide, *l*-Lysine *dl*-Methionine, *l*-Tryptophan *l*-Threonine, Salt, Ferrous Sulfate, Copper Sulfate, Choline Chloride, Manganous Oxide, *l*-Tryptophan, Niacin, Calcium Pantothenate, L-Carnitine, Copper Chloride, Vitamin A Supplement, Vitamin E Supplement, Menadione Dimethylpyrimidinol Bisulfite (source of K activity), Riboflavin Supplement, Vitamin B₁₂ Supplement, Thiamin, Vitamin D₃ Supplement, Ethylenediamine Dihydriodide, Sodium Selenite, Dried *Aspergillus oryzae* Fermentation Extract, Soybean Oil.

FEEDING DIRECTIONS

Full feed continuously to nursery pigs.

➔ WARNING ◀

Withdraw 2 days before to slaughter.

➔ CAUTION ◀

Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

➔ SPECIAL CONSIDERATIONS ◀

Protect medicated feed from excessive temperatures. Do not store at warm temperatures, 104° F (40°C), for more than 7 days.

➔ LIMITATIONS FOR USE ◀

Feed continuously as sole ration for 14 days. Not for use in swine weighing more than 250 lbs. Feed approximately 400 g/ton Chlortetracycline varying with body weight and feed consumption to provide 10 mg/lb of body weight per day.

Manufactured For
The Elite Swine Program, Inc
31822 St HWY CC
Hamilton, MO 64644

Net Weight 50 Lbs. (22.72 kg)