

# ELITE SWINE PRODUCT PIG PHASE 2 DEN/CTC MEDICATED

A COMPLETE PELLETTED FEED FOR NURSERY PIGS WEIGHING  
FROM LESS THAN 12 LB TO 15 LB.

Treatment of bacterial enteritis caused by Escherichia coli  
Salmonella choleraesuis and bacterial pneumonia caused by  
Pasteurella multocida susceptible to Chloretracycline. For the  
control of swine dysentery associated with Brachyspira  
(formerly Serpulina or Treponema) hyodsenteriae susceptible  
to Tiamulin.

## ACTIVE DRUG INGREDIENTS

Chlortetracycline.....	400 g/ton
Tiamulin.....	35 g/ton

## Guaranteed Analysis

Crude Protein.....Not Less Than.....	22.0%
Lysine..... Not Less Than.....	1.35%
Crude Fat.....Not Less Than .....	3.5 %
Crude Fiber .....	Not More Than..... 4.0%
Calcium (Ca).....Not Less Than.....	0.70%
Calcium (Ca).....Not More Than.....	1.20%
Phosphorus(P)..... Not Less Than.....	0.56%
Salt (NaCl).....Not Less Than.....	0.65%
Salt (NaCl).....Not More Than.....	1.15%
Selenium(Se).....Not Less Than.....	0.29ppm
Zinc(Zn).....Not Less Than.....	2000ppm

## INGREDIENTS:

Corn,Soybean Meal,Oats,Whey,Enzymatically Hydrolyzed Animal  
Protein (No Ruminant Proteins),Egg Product,Dried Lactobacillus  
acidophilus Fermentation Product,Grain By-Product Ethyl Alcohol,  
Ethyl Acetate Ammonium Poly Phosphate,Monocalcium Phosphate,  
Calcium Carbonate,Zinc Oxide, L-Lysine dl-Methionine,  
L-Tryptophan L-Theonine,Salt,Ferrous Sulfate,Copper  
Sulfate,Choline Chloride,Manganous Oxide,Niacin,Calcium  
Pantothenate,L-Carnitine,Copper Chloride,Vitamin A Supplement,  
Vitamin E Supplement,Menadione Dimethylpyrimidinol Bisulfite  
(source of K activity),Riboflavin Supplement,Vitamin B12  
Supplement,Thiamin,Vitamin D3 Supplement, Ethylenediamine  
Dihydrochloride,Sodium Selenite,Dried Aspergillus  
Fermentation Extract,Soybean oil.

## FEEDING DIRECTIONS:

Full feed continuously to nursery pigs.

**\*\*\* WARNING\*\*\***

Withdrawl 2 days before slaughter.

**\*\*\*\*CAUTION\*\*\***

Do not feed to animals other than swine. Use as the only source  
of Tiamulin and Chlortetracycline. Swine being treated with  
Tiamulin should not have access to feeds containing polyether  
ionophores(e.g. Monensin,Lasalocid,Narasin,Salinomycin and  
Semduramycin) as adverse reactions may occur. If signs of  
toxicity occur,discontinue use of medicated feeds.

**\*\*\* SPECIAL CONSIDERATIONS\*\*\***

Protect medicated feed from excessive temperatures. Do not  
store at warm temperature, 104 degree F(40 C) for more than 7  
days.

**\*\*\*LIMITATIONS FOR USE\*\*\***

Feed continuously as sole ration for 14 days. Not for use in  
swine weighing more than 250 lbs. Feeding approximately 400  
g/ton Chlortetracycline vary with body weight and feed  
consumption to provide 10 mg/lb of body weight per day.

Manufactured for  
The Elite Swine Program, Inc  
31822 St HWY CC  
Hamilton, MO 64644  
Net Weight 50 Lbs. (22.72 kg)